



Effects of Source and Rates of Potassium Fertilizer on Yield Traits and Potassium Use Efficiency of Potato in a Kenyan Ferralsol

Anthony O. Oketch ^{a,b}, Jonah K. Ngeno ^{b*},
Lucas K. Ngode ^b, Josiah C. Chiveu ^b and Harrison Churu ^c

^a Yara East Africa Ltd., P.O. Box 50949-00200, Nairobi, Kenya.

^b Department of Seed, Crop and Horticultural Sciences, University of Eldoret, P.O. Box 1125-30100, Eldoret, Kenya.

^c Department of Soil Science, University of Eldoret, P.O. Box 1125-30100, Eldoret, Kenya.

Authors' contributions

This work was carried out in collaboration among all authors. Author AOO executed field experiments and collected data. Author JKN wrote the manuscript. Author LKN reviewed and edited the manuscript. Author JCC analysed the data and reviewed materials and methods section. Author HC designed the hand held cardboard for measuring potato sizes and managed literature searches. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJSSPN/2023/v9i4201

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/105860>

Original Research Article

Received: 07/07/2023

Accepted: 10/09/2023

Published: 30/09/2023

ABSTRACT

Aims: Potassium (K) is one of the nutrient elements taken up in large amounts by potatoes, yet there is little research output on K fertilization of the crop in Kenya, due to the assumption that its natural soil reserves are sufficient for the production of most crops. The study objective was to determine the influence of two K sources (muriate of potash – KCL and sulphate of potash –

*Corresponding author: Email: jonah.ngeno@uoeld.ac.ke;

K₂SO₄) on K use efficiency and tuber yield components of potatoes in one agro-ecological zone of Kenya.

Place and Duration of Study: The experimental site was the University of Eldoret farm in Kenya, located at latitude 0°574942' N and longitude 35°300772' E, during the middle - second rains season (S1) and middle rains season (S2) of 2020 and 2021 respectively.

Methodology: Treatments were: 0 kg K ha⁻¹ (control), 60,120,180 and 240 kg K ha⁻¹ in S1 and 0, 30, 60, 90, 120 and 180 kg K ha⁻¹ in S2, supplied in the form of KCl and K₂SO₄. Treatments were laid in a Randomized Complete Block Design in 2x5 and 2x6 factorial arrangements in S1 and S2 respectively. The potato variety planted was Destiny. Data collected included: Agronomic efficiency of K (AE_K), total tuber yield, two categories of seed yield (Size 1 tubers with diameters of 28- 45 mm; Size 2 tubers with 46 - 60mm diameter), total seed yield and ware yield (tubers with diameters over 60mm).

Results: During the drier S1, the application of 120 kg K ha⁻¹ in the form of K₂SO₄ significantly ($P < .001$) increased the yield of size 1 tubers, total seed yield and total tuber yield. The total tuber yield increment from this K rate was only 12.6 % of the varietal potential yield, and its AE_K was 0.042 tonnes of tubers kg⁻¹ K. Therefore its application may not be economically viable. During the moisture-sufficient S2, all K fertilizer treatments in the form of KCl and 90 kg K ha⁻¹ in the form of K₂SO₄, contributed a yield increase of 58.1 - 68.6% and slightly exceeded varietal yield potential. Additionally, 30 kg K ha⁻¹ of KCl gave the highest AE_K of 0.5 tonnes of tubers kg⁻¹ K and consistently enhanced the yield of all potato sizes, except size 1.

Conclusion: Under similar ecological conditions as S2, potato farmers can achieve the highest AE_K and potential tuber yield with the application of 30 kg K ha⁻¹ of KCl. However, more trials on K nutrition in potatoes need to be done in diverse environments.

Keywords: Potato; potassium sulphate; potassium chloride; seed potato.

1. INTRODUCTION

Potato is the fourth most important food crop after rice, wheat and maize [1]. It is commonly used for making chips and crisps and has many industrial applications such as the manufacture of starch. The annual world production volume is slightly over 376 million metric tonnes, and the world average productivity is 20.7 tonnes ha⁻¹ [3]. Potato yields in Africa are still low, with only six countries recording average yields above 20 tons ha⁻¹ [2]. In the Eastern Africa region, the productivity of potatoes is 10.4 tonnes ha⁻¹, while the average productivity in Kenya is 9.8 tons ha⁻¹ against a potential of 40 tonnes ha⁻¹ [2,3]. The yield gap of 30 tons ha⁻¹ in Kenya can be attributed to several factors such as moisture deficiency during the cropping season since most farmers produce the crop under rain-fed conditions [4]. Bacterial wilt disease and poor soil health have been named as the second and third causes of the wide yield gap in Kenya and other Sub-Saharan countries [5]. Low soil fertility which is a component of poor soil health, has been cited as a key driver of low potato yields in Kenya [6]. Nitrogen (N) and phosphorous (P) deficiency contribute the most to low soil fertility and low potato yields in Kenyan soils [7,8]. Potassium (K) was thought to be

sufficient for potato production in most soil types of Kenya [9], and has received little research attention. However, in a study done in the major potato-growing areas of central and Eastern Kenya, 15-22% of farms sampled had K levels below the critical value of 200 mg kg⁻¹ soil [8]. Moreover, potato is known to take up more K than N and P for every ton of tuber produced [10]. In support of our hypothesis that K was likely to enhance tuber yields of potatoes, a supply of 45 kg K ha⁻¹ and optimal N, P and soil moisture yielded 63 tonnes ha⁻¹ of tubers in Andosols of Kenya [4]. Nonetheless, there is a knowledge gap on the choice of K source/type and fertilizer rates for tuber yield improvement in various soil types and seasons in Kenya. The results of this study will partly fill this information gap. The study was done in the ferralsols of Uasin-Gishu County, whose staple food crop for the past 50 years has been maize, and farmers are slowly adopting potato farming for food and income security. This is due to its short maturity period and a potential yield higher than three times that of maize. The objective of this study was to evaluate the effects of K source and rates on yield and K use efficiency of potato under rainfed conditions in the lower highland zone 3 of Uasin-Gishu county of Kenya.

2. MATERIALS AND METHODS

2.1 Site Description

The experiment was conducted at the University of Eldoret farm, located at a latitude of 0°57'49.42" N and longitude 35°30'07.72" E, in lower highland 3 (LH3) agro-ecological zone. The site has a nearly unimodal rainfall pattern, which is subdivided into three: First rainy season (end of March – end of June), middle rains (end of June to October) and second rains (November to February). The site is located at an altitude of 2135m above sea level, and receives mean annual rainfall and temperature of 1020mm and 17.3°C, respectively [11]. The initial soil analysis showed that the site has a pH (H₂O) of 5.71, total N of 0.18%, available P of 17.60 ppm, exchangeable K of 448.50 ppm, and organic carbon of 2.04%. The soil textural class is sandy clay. Soil analyses were done at the University of Eldoret soil laboratory following the procedures recorded in: [12]. During the middle-second rains season (September-December of 2020) (S1), the total rainfall received was 300mm, with the tuberisation to senescence growth stages receiving an average of 6.25 mm per week. During the middle rains season of 2021 (S2), the average seasonal rainfall was 645mm, and tuberisation to senescence growth stages received a mean of 50mm of rainfall per week; the mean temperature for S1 and S2 seasons were 17.4°C and 17.0°C respectively (Fig. 1).

2.2 Treatments, Experimental Design and Crop Management

The treatments for the S1 season were: 0 kg K ha⁻¹ (absolute control) 60, 120, 180 and 240 kg K ha⁻¹ in the form of KCL (muriate of potash - MOP) and K₂SO₄ (sulphate of potash - SOP). During the S2 season, the rates of K in the form of both KCL and K₂SO₄ were: 0 kg K ha⁻¹ (absolute control), 30, 60, 90, 120 and 180 kg K ha⁻¹. The additional rates during the second season were meant to give a better understanding of how K influences the parameters studied. K fertilizer was applied once during planting. Nitrogen fertilizer, (173.9 kg ha⁻¹) was applied in two splits in the form of urea, half rate at planting and half rate during the second earthing-up (fourth week after crop emergence). Phosphorus, 224.9 kg P₂O₅ ha⁻¹ was applied in the form of triple super phosphate at planting. Destiny potato variety obtained from AGRICO

limited was subjected to the treatments above. Planting furrows (about 23 cm deep) were made, and then potato seeds were placed in the furrows with no direct contact with fertilizer and then covered completely with soil. The treatments were laid out in a randomized complete block design (RCBD) in a 2x5 factorial arrangement in S1 and a 2x6 factorial arrangement in S2. Treatments were replicated three times. The plot size was 4.5 m x 4.5 m and crop spacing was 75 cm x 30 cm, giving a total of six crop rows. Weeding and earthing up of the crop was done manually using a hand hoe, during the second and fourth week after crop emergence. The crop was sprayed on a fortnightly basis using famoxadone-cymoxanil and acetamiprid, for crop protection against fungal diseases and insect pests. Crop harvesting was done carefully using a hand hoe, at harvest maturity - indicated by complete senescence and drying of crop shoots.

2.3 Data Collection

Harvesting of potato tubers was done on four inner rows, and the harvested tubers were graded based on sizes, before weighing of tubers as per the sizes using a digital weighing scale in kilograms (kg). Grading of potatoes into various sizes was done in each plot using an improvised hand-held measuring cardboard with holes of different diameters that correspond to potato sizes stipulated in Kenya Plant Health Inspectorate Service (KEPHIS) manual [13]. The sizes are: chatts (tubers with a diameter of less than 28mm); Size 1 (tubers with diameter between 28mm - 45mm); Size 2 (tubers with diameter between 46mm - 60mm) and ware (tubers with a diameter over 60mm). Sizes 1 and 2 are sold to the farmers for seed and ware is sold for culinary or industrial processing. Chatts are often utilised as animal feed. Total seed weight was obtained by adding weights of size 1 and 2 tubers in each plot; total tuber weight was obtained by the sum of the weights of chatts, size 1, size 2 and ware potato tubers in each plot. Tuber weights of each potato size were converted to tonnes ha⁻¹ based on the harvested area. The agronomic efficiency of applied potassium fertilizer (AE_K) was determined by: $AE_K = (Y - Y_0)/F$, where Y is the tuber yield with the application of K fertilizer, Y₀ is tuber yield without K fertilizer application and F is the amount of K applied [14].




Fig. 1. Mean monthly rainfall (A) and temperature (B) during the middle-second rains season of 2020 and middle rains season of 2021

2.3.1 Data analysis

Data collected was subjected to analysis of variance (ANOVA) using GenStat software 16th Edition (VSN International Ltd) at a 5% level of significance. Means were compared using Tukey's test.

3. RESULTS AND DISCUSSION

3.1 Results

It was observed that the application of 120 kg K ha⁻¹ in the form of SOP significantly increased the yield of chats, size 1, total seed yield and total tuber yield during the September-December rain season of 2020 (Table 1). It was further noted that potatoes supplied with 120 kg K ha⁻¹ in the form of K₂SO₄ had 5 more tons of the total tuber yield than the control, which was 29.5% of the actual yield (17.1 tons) and 12.6% of the potential yield (40 tonnes ha⁻¹). The K rate also gave the highest agronomic efficiency of K (Table 1). The larger tubers (ware) required 180kg K ha⁻¹ in the form of SOP to obtain the highest yield of 4.04 tonnes ha⁻¹, a 2.14-ton increase from the control mean (Table 1). Application of 240 kg K ha⁻¹ of either MOP or SOP reduced the yield of ware potato by 23 and 21% respectively, and total tuber yield by 11.7 and 3% respectively compared to the means of the control (Table 1).

During the middle rainy season of 2021, the application of 60 and 180 kg K ha⁻¹ in the form of MOP and SOP respectively, gave similar but significantly higher yield of chats compared to

the control (Table 2). Plants supplied with 60 kg K ha⁻¹ of both SOP and MOP and 120 kg K ha⁻¹ in the form of MOP significantly enhanced the yield of size 1 potato tubers; the yield contribution from the three K rates was between 41- 63%. The highest increment was attributed to the higher K rate (Table 2). All K rates from 30 kg ha⁻¹ in the form of MOP and only 180 kg K ha⁻¹ of SOP gave similar but significant yield increase of size 2 potato tubers compared to the controls (Table 2). The highest size 2 tuber yield increase (55%) was recorded in plants treated with 30 kg K ha⁻¹ of MOP. Generally, all plants treated with MOP at K rates from 30 kg ha⁻¹ significantly enhanced the total seed yield of potato tuber, but the SOP form of K enhanced the same seed yield only at rates of 60 and 180 kg K ha⁻¹ (Table 2). However, the highest total seed yield increment due to K fertilization was 44.1%, recorded in plants supplied with 60 kg K ha⁻¹ in the form of MOP. In summary, all potato plants supplied with K fertilizer irrespective of the form, had significantly higher total potato tuber yield compared to the controls during the middle rainy season (Table 2). Further analysis showed that all K fertilizer treatments in the form of KCl and 90 kg K ha⁻¹ in the form of K₂SO₄, contributed a yield increase of 58.1 - 68.6% and exceeded the varietal yield potential of 40 tonnes ha⁻¹ (Table 2).

Generally, it was observed that MOP increased total tuber yield more than SOP in the middle rains season of 2021, but during the moisture depressed middle and second rains seasons of 2020, SOP fertilizer gave more tuber yield compared to MOP (Fig. 2).

Table 1. Effects of potassium (K) fertilizer source and rate on total yield and yield of different sizes of potato tubers in a field experiment conducted at LH3 agro-ecological zone of Uasin-Gishu County of Kenya, between September and December of 2020 (end of middle - second rain season)

K Source	K rate	Yield of potato tubers in tonnes ha ⁻¹					Ware	Total yield	AE _K (kg tuber kg ⁻¹ K)
		Chatts	Size 1	Size 2	Total seed yield†				
MOP	0 kg ha ⁻¹	0.28 _d	3.32 _c	6.48	9.80 _{cd}	2.95 _b	13.03 _{bcd}	-	
MOP	60 kg ha ⁻¹	0.29 _d	3.98 _{bc}	7.97	11.96 _{ab}	1.70 _{cde}	13.94 _{bc}	31.50	
MOP	120 kg ha ⁻¹	0.44 _{bcd}	3.40 _c	5.14	8.54 _d	3.18 _{ab}	12.15 _{bcd}	0.83	
MOP	180 kg ha ⁻¹	0.61 _b	3.64 _c	5.96	9.60 _{cd}	2.45 _{bcd}	12.66 _{bcd}	3.39	
MOP	240 kg ha ⁻¹	0.38 _{cd}	3.64 _c	5.16	8.80 _d	1.46 _{de}	10.64 _e	-5.90	
SOP	0 kg ha ⁻¹	0.28 _d	3.33 _c	6.61	9.93 _{cd}	0.85 _e	11.06 _{de}	-	
SOP	60 kg ha ⁻¹	0.56 _b	4.82 _b	6.08	10.90 _{bc}	2.28 _{bcd}	13.73 _{bc}	28.00	
SOP	120 kg ha ⁻¹	0.83 _a	6.02 _a	7.69	13.71 _a	2.56 _{bc}	17.10 _a	42.08	
SOP	180 kg ha ⁻¹	0.51 _{bc}	3.30 _c	6.31	9.61 _{cd}	4.04 _a	14.16 _b	11.72	
SOP	240 kg ha ⁻¹	0.47 _{bc}	4.04 _{bc}	5.64	9.68 _{cd}	1.50 _{cde}	11.66 _{cde}	-1.63	
SEM[§]		0.03	0.23	0.38	0.40	0.21	0.46		
P value		<.001	<.001	‡	<.001	<.001	<.001		

Means followed by similar subscript letters along a column are not significantly different ($P \leq 0.05$). † Sum of size 1&2, MOP- KCL, SOP- K₂SO₄. ‡- means similar to controls though P value was =.05. § Standard error of mean

Table 2. Effects of potassium (K) fertilizer source and rate on total yield and yield of different sizes of potato tubers in a field experiment conducted at LH3 agro-ecological zone of Uasin-Gishu County of Kenya, between June and October of 2021 (middle rainy season)

K Source	K rate	Yield of potato tubers in tonnes ha ⁻¹					Ware	Total yield	AE _K (kg tuber kg ⁻¹ K)
		Chatts	Size 1	Size 2	Total seed yield†				
MOP	0 kg ha ⁻¹	0.43 _{bcde}	3.78 _{cd}	9.99 _d	13.77 _e		11.75 _c	25.95 _c	-
MOP	30 kg ha ⁻¹	0.51 _{abcde}	3.79 _{cd}	15.57 _a	19.36 _a		21.18 _{ab}	41.05 _a	503
MOP	60 kg ha ⁻¹	0.65 _a	5.33 _{ab}	14.52 _{ab}	19.85 _a		22.59 _a	43.08 _a	286
MOP	90 kg ha ⁻¹	0.43 _{bcde}	4.94 _{abc}	14.20 _{abc}	19.14 _{ab}		23.67 _a	43.24 _a	192
MOP	120 kg ha ⁻¹	0.62 _{ab}	6.19 _a	13.22 _{abc}	19.42 _a		23.27 _a	43.31 _a	145
MOP	180 kg ha ⁻¹	0.31 _e	4.07 _{bcd}	13.70 _{abc}	17.77 _{abcd}		25.68 _a	43.77 _a	99
SOP	0 kg ha ⁻¹	0.43 _{bcde}	3.78 _{cd}	9.99 _d	13.77 _e		11.75 _c	25.95 _c	-
SOP	30 kg ha ⁻¹	0.37 _{de}	3.7 _{cd}	12.03 _{bcd}	15.73 _{cde}		16.27 _{bc}	32.37 _b	214
SOP	60 kg ha ⁻¹	0.56 _{abc}	5.73 _a	11.92 _{cd}	17.65 _{abcd}		14.24 _c	32.47 _b	109
SOP	90 kg ha ⁻¹	0.57 _{abcd}	3.98 _{bcd}	12.33 _{bcd}	16.32 _{bcde}		24.88 _a	41.77 _a	176
SOP	120 kg ha ⁻¹	0.38 _{cde}	3.09 _d	11.79 _{cd}	14.89 _{de}		16.23 _{bc}	31.49 _b	46
SOP	180 kg ha ⁻¹	0.71 _a	4.04 _{bcd}	14.23 _{abc}	18.26 _{abc}		13.12 _c	32.09 _b	34
SEM[§]		0.04	0.29	0.49	0.58		1.18	1.05	
P value		<.001	<.001	.003	.001		<.001	<.001	

Means followed by similar subscript letters along a column are not significantly different ($P \leq 0.05$). † Sum of size 1&2, MOP- KCL, SOP- K₂SO₄. § standard error of mean




Fig. 2. Effects of potassium source and season on total tuber yield of potato in a field experiment conducted at the University of Eldoret farm during the middle and second rains season of 2020 (September –December) and middle rains season of 2021 (June – October)

3.2 Discussion of Results

3.2.1 K source and season interactions

Application of KCL form of K fertilizer can lead to the attainment of potential tuber yield of potato in the semi-humid tropical soil when soil moisture is adequate for crop production. However, during periods of soil moisture deficiency, K_2SO_4 form of K can give better tuber yields compared to KCL. Previous researchers have reported a higher yield contribution of K_2SO_4 than KCL [15,16], but a similar yield increase from KCL than K_2SO_4 has been reported in one potato genotype [17]. A review of 48 papers on K nutrition in potato shows that there is no consistent superiority of either K_2SO_4 or KCL on tuber yield increment in potato [18]. The same authors observed that the response of potatoes to K may be influenced by genotype, moisture availability, and soil physiochemical conditions among others. In this study, higher yield response due to KCL may be attributed to sufficient moisture, which was within weekly recommended range between 12.7mm (at emergence) - 50mm (at tuberisation to senescence) (https://cropwatch.unl.edu /potato /plant_g rowth). Results from this study also confirm that certain potato varieties are less sensitive to Cl⁻ in KCL [19]. Higher yield increases due to K_2SO_4 than KCL in drier season may be attributed to drought stress tolerance conferred by potassium sulphate [20], possibly due to improvement of water use efficiency in potato [21]. However, the yields were generally lower than 50% of the potential yield during the drier season. This observation may have been caused by low moisture availability which declined from 25mm per week during emergence

to a mean of about 6.25 mm/week from the 45th day to senescence, a shortfall of over 40mm/week (https://cropwatch.unl.edu /potato /plant_g rowth). Therefore, the contribution of K_2SO_4 to yield enhancement of potato may not be economically viable during the moisture depressed season, owing to the higher fertilizer costs in Africa compared to the rest of the world [22].

3.2.2 K source and K rate interactions

Potassium sulphate application at the rate of 120 kg K ha⁻¹ enhanced 28 - 45mm sized tubers, total seed and total tubers yields during the drier season, but agronomic efficiency of K at this K rate was low, and this can explain why the contribution of the K rate to the potential yield was less than 13%. However, beyond 120 kg K ha⁻¹ of K_2SO_4 application, tuber yield started declining and negative yields relative to the control were recorded at 240 kg K ha⁻¹. This was a contradiction of earlier findings where the yield response of potato to K_2SO_4 was recorded up to 280 kg ha⁻¹ [23]. Despite the report that increasing K_2SO_4 fertiliser rates can enhance water stress tolerance by crops [24], the physiological explanation for the reduced potato yields with increasing K_2SO_4 fertiliser in this study, needs further investigation.

On the other hand, all K rates of KCL above 30 kg K ha⁻¹ in this study enhanced the yield of tubers above 45mm, total seed and total tuber yield during the middle rains. Although the yield increases were statistically similar at rates between 30-180 kg K ha⁻¹ of KCL, the agronomic efficiency of K was highest at 30 kg K ha⁻¹, which

indicates that the plants treated with higher K rates were taking up K luxuriously, with no significant yield contribution [25]. Most studies have reported higher yield response to K in the form of KCL at rates beyond 300 kg K ha⁻¹ [23], but the yield contribution of K at lower rates in this study may be attributed to higher levels of exchangeable K (448 mg kg⁻¹ soil). In a related study conducted in tropical acidic soils, potato yield response to 165 kg K ha⁻¹ in the form of KCL was attained when available soil K was up to 145 mg kg⁻¹ [26]. In Central highland soils of Kenya with available soil K of 156 mg kg⁻¹, a slight potato yield decline was observed when KCL fertilizer was applied at rates of 224 kg K ha⁻¹ [8]. It was an interesting observation to report yield increment due to K in this study, beyond the recommended critical soil K level of 200 kg ha⁻¹ [18]. This suggests that there might be other ecological factors that may influence potato response to K other than the available soil K.

4. CONCLUSION

Seasonal moisture variation is likely to be one of the key ecological factors influencing the response of potatoes to KCL or K₂SO₄ forms of K in tropical semi-humid soils that depend entirely on rainfall for crop production. K rate of 30 kg K ha⁻¹ supplied as KCL is sufficient for the production of potato tubers above 45mm in diameter in regions with similar soil conditions as the study area, and similar weather conditions recorded in the middle rainy season.

5. RECOMMENDATIONS OF THE STUDY

It is recommended that farmers relying on rainfall for potato production in similar agro-ecological zones as the study area can apply 30kg ha⁻¹ of KCL for optimal tuber yield of potato in seasons when soil moisture is sufficient. Detailed season-based studies need to be done to understand K fertilizer use efficiency in diverse soil types.

ACKNOWLEDGEMENTS

The authors wish to acknowledge the field and technical staff who assisted in data collection and soil analyses. We also acknowledge funding support from the University of Eldoret annual research grant – cohort 5.

COMPETING INTERESTS

Authors have declared that they have no known competing financial interests or non-financial interests or personal relationships that could

have appeared to influence the work reported in this paper.

REFERENCES

1. Zhang H, Xu F, Wu Y, Hu H-h, Dai X-f. Progress of potato staple food research and industry development in China. J Integr Agric. 2017;16(12):2924-32.
2. Devaux A, Goffart J-P, Kromann P, Andrade-Piedra J, Polar V, Hareau G. The potato of the future: Opportunities and challenges in sustainable agri-food systems. Potato Res. 2021;64(4):681-720.
3. FAO STAT. Crops and livestock products. 2023. Accessed 12 August; 2023. Available: <https://www.fao.org/faostat/en/#data/QCL>
4. Satognon F, Owido SFO, Lelei JJ. Effects of supplemental irrigation on yield, water use efficiency and nitrogen use efficiency of potato grown in mollic Andosols. Environ Syst Res. 2021;10(1):38.
5. Harahagazwe D, Condori B, Barreda C, Bararyenya A, Byarugaba AA, Kude DA, et al. How big is the potato (*Solanum tuberosum* L.) yield gap in sub-saharan africa and why? A participatory approach. Open Agric. 2018;3(1):180-9
6. Muthoni J. Soil fertility situation in potato producing kenyan highlands—case of KALRO-Tigoni. Int. J. Hortic. 2016;6(25):1-11.
7. NAAIAP. Soil suitability evaluation for maize production in Kenya. Nairobi: ministry of agriculture, livestock and fisheries; 2014.
8. Mugo JN, Karanja NN, Gachene CK, Dittert K, Nyawade SO, Schulte-Geldermann E. Assessment of soil fertility and potato crop nutrient status in Central and Eastern highlands of Kenya. Sci Rep. 2020;10(1):7779.
9. Recke H, Schnier HF, Nabwile S, Qureshi JN. Responses of Irish potatoes (*Solanum tuberosum* L.) to mineral and organic fertilizer in various agro-ecological environments in Kenya. Exp Agric. 1997;33(1):91-102.
10. YARA. Crop nutrition: potatoes. 2019. Accessed 17 August 2023. Available:<https://www.yara.co.ke/crop-nutrition/potato/potato-nutritional-summary/>.
11. Jaetzold R, Schmidt H, Shisanya C. Farm management handbook of Kenya vol. II, subpart B1b, Uasin-Gishu County.

- Nairobi:Ministry of Agriculture and GIZ; 2011.
12. Okalebo JR, Gathua KW, Woomer PL. Laboratory methods of soil and plant analysis: A working manual. 2nd ed. TSBF-CIAT and SACRED Africa, Kenya; 2002.
 13. Kephis. Seed potato production and certification guidelines. 2016. Accessed 1st August 2023. Available:<https://www.kephis.org/images/docs/seedpotatobooklet.pdf>.
 14. Fixen P, Brentrup F, Bruulsema T, Garcia F, Norton R, Zingore S. Nutrient/fertilizer use efficiency: Measurement, current situation and trends. In: Pay Drechsel, Patrick Heffer, Hillel Magen, Robert Mikkelsen, Wichelns D, editors. Managing water and fertilizer for sustainable agricultural intensification: International Fertilizer Industry Association (IFA), International Water Management Institute (IWMI), International Plant Nutrition Institute (IPNI), International Potash Institute (IPI); 2015.
 15. Mello SdC, Pierce FJ, Tonhati R, Almeida GS, Neto DD, Pavuluri K. Potato response to polyhalite as a potassium source fertilizer in Brazil: yield and quality. HortSci 2018;53(3):373-9.
 16. Geremew T, Ayalew A, Getachew. Response of potato (*Solanum tuberosum* L.) to potassium fertilizer on acid soils of Wolmera and Gumer Weredas, in the highlands of Ethiopia. J Biol Agric Healthc. 2015;5(17):156-60.
 17. Wilmer L, Pawelzik E, Naumann M. Comparison of the effects of potassium sulphate and potassium chloride fertilisation on quality parameters, including volatile compounds, of potato tubers after harvest and storage. Front Plant Sci. 2022;13:920212.
 18. Torabian S, Farhangi-Abriz S, Qin R, Noulas C, Sathuvalli V, Charlton B, et al. Potassium: A vital macronutrient in potato production—A review. Agron. 2021; 11(3):543.
 19. Koch MT, Pawelzik E, Kautz T. Chloride changes soil–plant water relations in potato (*Solanum tuberosum* L.). Agron. 2021;11(4):736.
 20. Bahar AA, Faried HN, Razzaq K, Ullah S, Akhtar G, Amin M, et al. Potassium-induced drought tolerance of potato by improving morpho-physiological and biochemical attributes. Agron. 2021; 11(12):2573
 21. Darwish T, Fadel A, Chahine S, Baydoun S, Jomaa I, Atallah T. Effect of potassium supply and water stress on potato drought tolerance and water productivity. Commun Soil Sci Plant Anal. 2022;53(9):1100-12.
 22. Kitonyo OM, Sadras VO, Zhou Y, Denton MD. Nitrogen fertilization modifies maize yield response to tillage and stubble in a sub-humid tropical environment. Field Crops Res. 2018;223:113-24.
 23. Panique E, Kelling KA, Schulte EE, Hero DE, Stevenson WR, James RV. Potassium rate and source effects on potato yield, quality, and disease interaction. Am Potato J. 1997;74(6): 379-98.
 24. Mohd Zain, Nurul Amalinalsmail, Razi M. Effects of potassium rates and types on growth, leaf gas exchange and biochemical changes in rice (*Oryza sativa*) planted under cyclic water stress. Agric Water Manag. 2016;164:83-90.
 25. Kang W, Fan M, Ma Z, Shi X, Zheng H. Luxury absorption of potassium by potato plants. Am J Potato Res. 2014;91: 573-8.
 26. Job LG, Soratto RP, Fernandes AM, Assunção NS, Fernandes FM, Yagi R. Potassium fertilization for fresh market potato production in tropical soils. Agron J. 2019;111(6):3351-62.

© 2023 Oketch et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/105860>